How to take your medicine

Your doctor has prescribed this medicine (Veltassa®) for you to help lower the levels of potassium in your blood.

It is easy to take – just follow the 3 steps below.





Mix 3 tablespoons (40 mL) of water with an entire sachet of Veltassa®, then stir.

2. ADD



Add another 3 tablespoons (40 mL) of water and stir thoroughly. The powder will not dissolve and the mixture will look cloudy.*

3. DRINK



Drink immediately. If powder remains in the glass after drinking, repeat stages 2 and 3 to ensure the entire dose is taken.

Model has been used to represent a patient

You need to take the medicine **once a day with or without food**. You should not heat your medicine or add it to heated foods or liquids. You can store it below 25°C for up to 6 months.

It is important to leave at least a 3-hour gap between taking this and other medicines. For example, if you need to take another medicine with your lunch at 13.00 you can take this medicine with an afternoon snack at 16:00. Any other medicines can then be taken after 19.00.

To help keep track of when you have taken your medicines please print this page. The section below has space for you to note down when you have taken your medicines.

As with any medicine, you should always read the patient information leaflet. If you have any queries or concerns, please talk to your doctor.

*You may add more water to the mixture to help you swallow the medicine. If preferred you can mix Veltassa® with apple or cranberry juice.



The information and materials for Veltassa® contained in this website were prepared based on the EU SmPC. Prescribing information may vary depending on local approval in each country.

Therefore, before prescribing any product, always refer to local materials such as the prescribing information and/or the Summary of Product Characteristics (SPC).





Date of preparation: November 2020 | HQ-PAT-2000065